EN530.678 Nonlinear Control and Planning in Robotics Homework #1

February 7, 2022

Due: February 14, 2022 (before class)

Prof: Marin Kobilarov

- 1. Perform the following calculations (2 pts each):
 - (a) Let $f(x) = ||x||^3$ where $x \in \mathbb{R}^n$. Compute the gradient $\nabla f(x)$ and Hessian $\nabla^2 f(x)$, ideally using vector calculus as opposed to individual coordinates x_i .
 - (b) Let $\dot{x} = Ax + b$ where $x \in \mathbb{R}^n$ and V(x) = ||x||. What is $\dot{V}(x)$?
 - (c) Show that $||a \times b||^2 = ||a||^2 ||b||^2 (a^T b)^2$ for some $a, b \in \mathbb{R}^3$
 - (d) Show that $a \times (b \times c) = (a^T c)b (a^T b)c$ for some $a, b, c \in \mathbb{R}^3$
- 2. (Khalil) For each of the following systems, use a quadratic Lyapunov function candidate (e.g. $V(x) = \frac{1}{2}x^T x$ or more generally $V(x) = x^T A x$ for some matrix A > 0) to show that the origin is asymptotically stable. (3 pts each)
 - (a) $\dot{x}_1 = -2x_1$, $\dot{x}_2 = 2x_1 x_2$ (b) $\dot{x}_1 = -x_1(1+x_2^2)(1-x_1^2-x_2^2)$, $\dot{x}_2 = -x_2(1-x_1^2)(1-x_1^2-x_2^2)$ (c) $\dot{x}_1 = -(x_1+x_2)(1+2x_1^2-4x_1^4)$, $\dot{x}_2 = x_1(1+2x_1^2-4x_1^4)$ (d) $\dot{x}_1 = -x_1 - 5x_1x_2^2$, $\dot{x}_2 = 2x_1^2x_2 - x_2^3$
- 3. Implementation: For 2(a)-2(d), write a Matlab or Python script which plots x(t) and your Lyapunov candidate V(t) for $t \in [0, 5]$ starting from x(0) = (0.5, 0.5). You can use the ode45 function for integrating the dynamics. An example script is provided on the course web page for reference. See file hw1_lyapunov_example.m. (2 pts each)

Note: Upload your code and plots as a .zip file using https://forms.gle/Z2AYx3FRNJHtXTR47; in addition attach a printout of the code and all plots to your homework solutions.

4. (Khalil) Consider the system

$$\dot{x} = -a[I_n + S(x) + xx^T]x,$$

where a is a positive constant, I_n is the nxn identity matrix, and S(x) is an x-dependent skew symmetric matrix. Show that the origin is globally asymptotically stable. (10 pts)